



Curevita Research  
<https://Curevita.org>

Contents lists available at [Curevita Journals](https://www.curevitajournals.com)

## Frontiers of Agri and animal Innovation

journal homepage: [www.curevitajournals.com](https://www.curevitajournals.com)



# To evaluate the effect of strategic nutrient supplementation on the growth performance of Murrah buffalo calves

**Neeha Meena, Sandeep Nanavati, Mohabbat Singh Jamra, Ashok Kumar Patil, Nawal Singh Rawat, Danveer Singh Yadav,**

1. Livestock production and Management, College of Veterinary Science and A.H.,(NDVSU), Mhow, Indore, M.P.
2. Department of Animal Nutrition, College of Veterinary Science and A.H., (NDVSU), Mhow, Indore, M.P.

### Articalinfo

**Article history:** Received 22 June 2025, Revised 18 Aug 2025, Accepted 20 Aug 2025, Published Sept 2025

**Keywords:** Strategic supplementation, Nutrients, Growth performance, Murrah buffalo, Calves.

**Corresponding Author:** Neeha Meena, Department of Livestock Production and Management, College of Veterinary Science and A.H., (NDVSU), Mhow, Indore, (M.P.), India.

**Email:** [meenaneha602@gmail.com](mailto:meenaneha602@gmail.com)

**Citation:** Meena Neeha, Nanavati Sandeep, Jamra Mohabbat Singh Jamra, Patil Ashok Kumar, Rawat Nawal Singh, and Yadav Danveer Singh. 2025. To evaluate the effect of strategic nutrient supplementation on the growth performance of Murrah buffalo calves. Frontiers of Agri & Animal Innovation. 1,1,19-44..

**Publisher:** Curevita Research Pvt Ltd

### Abstract

A total of eighteen Murrah buffalo calves, aged 6-9 months and with similar body weights, were randomly selected and assigned to three equal groups (n=6). In control group ( $T_0$ ) animals were maintained on common feeding regimen consisting of required concentrate mixture and roughage as per requirement given by farmer during survey, in treatment group ( $T_1$ ) was given basal ration and strategic macronutrients (based on deficiency of macronutrients observed during survey) and in treatment group ( $T_2$ ) was given basal ration and strategic macronutrients and strategic micronutrients (based on deficiency of micronutrients observed during survey) for 90 days.



## Indtroduction

Body weight changes, feed intake and average daily gain were recorded weekly to assess growth performance for 90 days of the experiment. Fortnightly body weights were measured using an electronic weighing balance to evaluate the animals' weight. The overall average body weight was  $73.74 \pm 6.27$  for the T<sub>0</sub> group,  $79.2 \pm 4.9$  for the T<sub>1</sub> group, and  $81.67 \pm 2.5$  for the T<sub>2</sub> group, with no statistically significant differences ( $p > 0.05$ ) detected. However, numerically, growth was 7.43 and 10.75 percent higher in the T<sub>1</sub> and T<sub>2</sub> groups, respectively. The overall average daily gain was  $247.08 \pm 04.75$  for the T<sub>0</sub> group,  $418.17 \pm 9.4$  for T<sub>1</sub> group and  $465.2 \pm 6.81$  for the T<sub>2</sub> group, with a statistically significant difference. The average feed conversion efficiency was  $13.15 \pm 0.32$  for T<sub>0</sub> group,  $8.22 \pm 0.2$  for T<sub>1</sub>

group and  $7.6 \pm 0.15$  for the T<sub>2</sub> group, with significant differences observed. The p-value for the feed conversion ratio was ( $p \leq 0.01$ ).

### Formulation of a strategic nutrient supplement for Murrah buffalo calves

Based on the results obtained in phase I, the availability of crude protein and TDN of calves was compared with the standard nutrient requirements of calves by ICAR (2013). There was a deficiency of CP and TDN to the tune of 79.17% and 49.86%, respectively. For the preparation of a strategic mineral mixture supplement, zinc sulphate, cobalt chloride, and sodium dihydrogen ortho phosphate dihydrate were used as per the requirements given by ICAR (2013) as shown in Table -1. The measured quantity of strategic mineral mixture was supplemented in the ration of calves of the treatment group. This strategic mineral mixture was then added in the ratio of six Murrah



buffalo calves and a feeding trial was conducted.

**Experimental animals:** Eighteen Murrah buffalo calves of approximately similar age (6-9 months) and body weight as shown in Table-2, were selected and randomly allotted into three equal groups (6 in each group). All calves were maintained on a common feeding regime consisting of required concentrate mixture, wheat straw and green as per requirements given by ICAR (2013).

#### **Housing and management**

All the experimental calves were housed in a well-ventilated shed with provision of individual feeding and watering. Strict management and hygienic practices were adopted throughout the experimental period. Clean drinking water was provided

*ad libitum* twice a day at about 10:00 a.m. and 5:00 p.m. daily.

#### **Feeds and feeding**

The calves were offered wheat straw, green and concentrate ration to meet their nutrient requirements in Table 03 recommendation for a body weight gain of 70kg/day. All the experimental calves were fed on a basal diet comprised of a concentrate ration, green and wheat straw. Weighed amount of concentrate ration was provided at 10 a.m. daily to meet *almost* their whole CP and a major part of TDN requirement. Wheat straw was provided *ad libitum* after complete consumption of the concentrate mixture. The amount of concentrate mixture required by each calf was adjusted at every fifteen-day interval based on their body weights.



**Table-1: Composition of strategic micronutrient supplementation (animal/day) for Murrah buffalo calves**

Composition	Quantity (mg)	P (mg)	Co (mg)	Zn (mg)
Zinc sulphate (363.6 mg/g)	36.3	-	-	13.2
Cobalt chloride (250 mg/g)	1.92	-	0.48	-
Sodium dihydrogen ortho phosphate dihydrate (172.5 mg/g)	10.72	1.85	-	-
Total supply		1.85	0.48	13.2

**Table-2: Initial body weight (kg) of Murrah buffalo calves**

S.No.	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>
1	51.25	52.9	57.60
2	52.13	55.75	58.45
3	61.07	56.45	67.75
4	72.25	77.05	65.15
5	88.17	77.35	70.65
6	51.85	53.85	56.35
Overall mean $\pm$ SE	62.79 $\pm$ 6.06	62.23 $\pm$ 4.76	62.66 $\pm$ 2.44

\

**Table-3: The feeding schedule of experimental Murrah buffalo calves**

Groups	No. of Calves	Feeding on	Period (days)
T <sub>0</sub>	06	Basal Ration (Wheat straw+Green+Concentrate mixture)	90
T <sub>1</sub>	06	Basal Ration + strategic macronutrients (based on deficiency of macronutrients observed during survey)	90
T <sub>2</sub>	06	Basal Ration + strategic macronutrients + strategic micronutrients (based on deficiency of micronutrients observed during survey)	90

**Recording of body weight, dry matter intake and feed conversion ratio**



The body weight of all the experimental animals was recorded at fortnightly intervals during the morning time before offering them any feed and water. The measurement was made in kilograms by individually weighing the calves using an electronic weighing balance. Calves were offered a weighed

amount of concentrate mixture, green and gram straw. Residue, if any was weighed after next day morning. Samples of concentrate mixture, green and gram straw were subjected for DM analysis once in a fortnight, to know their DM intake. The feed conversion ratio (Banerjee,1998) was calculated using the formula:

$$FCR = \frac{\text{Feed intake (g)}}{\text{Body weight gain (g)}}$$

Data were analyzed using one-way ANOVA and Duncan's multiple range test in SPSS 20.0.(Snedecor and Cochran, 1994)

## Results and Discussion

### Chemical composition of feedstuffs (% DM basis)

The average percentages of proximate components, including crude protein (CP), ether extract (EE), crude fiber (CF), total ash (TA), and nitrogen-free extract (NFE) on a dry matter (DM) basis, for the available feedstuffs are detailed in Table 04.

The available feeds are wheat straw, gram straw, masoor straw, Green (Local grass), Green maize, compound feed, cottonseed cake and Wheat bran.

Among the straws, DM content of wheat straw was  $92.83 \pm 0.20$  % and the crude protein (CP), ether extract (EE), crude fibre (CF), total ash (TA) and nitrogen-free extract (NFE) contents were  $2.65 \pm 0.28$  %,  $1.35 \pm 0.14$  %,  $37.2 \pm 0.61$  %,  $06.52 \pm 0.22$  % and  $52.29 \pm 1.19$  %, respectively. Similarly, the DM content of gram



straw was  $91.38 \pm 0.19$  % and the CP, EE, CF, TA and NFE contents were  $8.42 \pm 0.21$  %,  $1.27 \pm 0.20$  %,  $38.22 \pm 0.17$  %,  $11.50 \pm 0.25$  % and  $40.58 \pm 0.25$  %, respectively. The DM content of masoor straw was  $93.62 \pm 0.18$  % and the CP, EE, CF, TA and NFE contents were  $6.47 \pm 0.26$  %,  $2.37 \pm 0.22$  %,  $39.30 \pm 0.17$  %,  $09.43 \pm 0.28$  % and  $42.43 \pm 0.23$  %, respectively.

Among the green roughages, DM content of local grass was  $25.87 \pm 0.62$  % and the CP, EE, CF, TA and NFE contents were  $6.53 \pm 0.22$  %,  $1.39 \pm 0.17$  %,  $34.63 \pm 0.33$  %,  $09.40 \pm 0.21$  % and  $48.04 \pm 0.63$  %, respectively. Similarly, the DM content of green maize was  $20.38 \pm 0.19$  % and the CP, EE, CF, TA and NFE contents were  $6.12 \pm 0.12$  %,  $1.35 \pm 0.22$  %,  $30.27 \pm 0.30$  %,  $11.81 \pm 0.18$  % and  $50.45 \pm 0.26$  %, respectively.

Among the concentrate feeds, DM content of compound feed was  $91.93 \pm 0.43$  % and the CP, EE, CF, TA and NFE contents were  $18.73 \pm 0.23$  %,  $2.45 \pm 0.26$  %,  $11.95 \pm 0.36$  %,  $10.7 \pm 0.32$  % and  $56.17 \pm 0.41$  %, respectively. Similarly, the DM content of cotton seed cake was  $91.12 \pm 0.46$  % and the CP, EE, CF, TA and NFE contents were %,  $24.93 \pm 0.07$  %,  $3.32 \pm 0.18$  %,  $22.57 \pm 0.35$  %,  $04.80 \pm 0.04$  % and  $44.39 \pm 0.16$  %, respectively. The DM content of wheat bran was  $92.37 \pm 0.21$  % and the CP, EE, CF, TA and NFE contents were %,  $13.26 \pm 0.17$  %,  $3.45 \pm 0.21$  %,  $12.37 \pm 0.16$  %,  $04.58 \pm 0.18$  % and  $66.35 \pm 0.13$  %, respectively.

**Table 04: Chemical composition of feedstuffs (% DM basis)**

Sample	DM (%)	CP (%)	EE (%)	CF (%)	TA (%)	NFE (%)
Wheat Straw	92.83 ± 0.20	2.65 ± 0.28	1.35 ± 0.14	37.2 ± 0.61	06.52 ± 0.22	52.29 ± 1.19
Gram straw	91.38 ± 0.19	8.42 ± 0.21	1.27 ± 0.20	38.22 ± 0.17	11.50 ± 0.25	40.58 ± 0.25
Masoor straw	93.62 ± 0.18	6.47 ± 0.26	2.37 ± 0.22	39.30 ± 0.17	09.43 ± 0.28	42.43 ± 0.23
Green (Local grass)	25.87 ± 0.62	6.53 ± 0.22	1.39 ± 0.17	34.63 ± 0.33	09.40 ± 0.21	48.04 ± 0.63
Green maize	20.38 ± 0.19	6.12 ± 0.12	1.35 ± 0.22	30.27 ± 0.30	11.81 ± 0.18	50.45 ± 0.26
Compound feed	91.93 ± 0.43	18.73 ± 0.23	2.45 ± 0.26	11.95 ± 0.36	10.7 ± 0.32	56.17 ± 0.41
Cotton seed cake	91.12 ± 0.46	24.93 ± 0.07	3.32 ± 0.18	22.57 ± 0.35	04.80 ± 0.04	44.39 ± 0.16
Wheat bran	92.37 ± 0.21	13.26 ± 0.17	3.45 ± 0.21	12.37 ± 0.16	04.58 ± 0.18	66.35 ± 0.13

**Average mineral content of feedstuffs (% DM basis)**

The mean mineral content in feedstuffs is given in Table 05. Among straws, wheat straw had the highest P ( $0.15 \pm 0.01$  %) and Mn ( $43.23 \pm 2.5$ ) whereas lowest Ca ( $0.45 \pm 0.03$  %), and Co ( $0.07 \pm 0.00$  ppm), while gram straw had the highest Ca ( $1.46 \pm 0.16$  %), Cu ( $8.35 \pm 0.18$  ppm) and Co ( $0.63 \pm 0.07$  ppm) and lowest P ( $0.11 \pm 0.02$  %), Fe ( $360.63 \pm 1.37$  ppm), Zn ( $5.26 \pm 0.17$  ppm) and Mn ( $35.45 \pm 0.28$ ). Masoor straw was found to have the highest Fe ( $606.63 \pm 2.998$  ppm) and Zn ( $23.04 \pm 0.05$  ppm), there after lowest Cu ( $4.48 \pm 0.14$  ppm). In concentrate feeds, cotton seed cake had the lowest Cu ( $8.68 \pm 0.35$  ppm) and Mn ( $32.28 \pm 0.25$  ppm) while highest Co ( $0.53 \pm 0.013$  ppm). Wheat bran had the highest Zn ( $62.07 \pm 0.17$  ppm) and Mn ( $130.68 \pm 0.36$  ppm) but the lowest Co ( $0.2 \pm 0.05$  ppm) and Ca ( $0.22 \pm 0.05$  ppm). The compound feed was notable for the highest P ( $0.81 \pm 0.05$  ppm), Fe

( $744.9 \pm 0.82$  ppm), Cu ( $18.53 \pm 1.07$  ppm) and Ca ( $0.91 \pm 0.05$  ppm), while the lowest Zn ( $20.23 \pm 1.17$  ppm). Among green roughages, green maize fodder had the highest Ca ( $0.55 \pm 0.13$  %), Mn ( $130.73 \pm 1.34$  ppm) and Cu ( $8.33 \pm 0.14$  ppm) while local grasses were found to have higher Zn ( $12.23 \pm 0.71$  ppm) and P ( $0.17 \pm 0.01$  ppm) compared to other green feeds.

**Nutritional status of Murrah buffalo calves body weight and feed intake**

The body weight and dry matter (DM) intake (kg/day) from different feed ingredients for Murrah buffalo calves are displayed in Table 05. The average body weight of the calves was  $64.16 \pm 1.35$  kg. The average total DM intake was  $1.38 \pm 0.01$  kg/day. The average DM intake (kg/day) from wheat straw, green forage and concentrate was  $1.04 \pm 0.01$ ,  $0.20 \pm 0.01$ , and  $0.15 \pm 0.01$ , respectively.





**Table-5: Dry matter intake (kg/d) of different feed ingredients offered to Murrah buffalo calves**

Animal no.	Body weight (kg)	Wheat straw (kg)	Green (kg)	Concentrate(kg)	Total DMI (kg)
1	59.95	1.03	0.17	0.13	1.33
2	62.53	1.00	0.16	0.17	1.33
3	61.67	1.00	0.19	0.11	1.30
4	62.75	1.06	0.13	0.15	1.34
5	58.87	1.05	0.25	0.17	1.47
6	61.75	1.05	0.18	0.16	1.39
7	62.90	1.03	0.19	0.12	1.34
8	65.75	1.05	0.17	0.13	1.35
9	55.87	1.02	0.20	0.17	1.39
10	65.85	1.08	0.24	0.12	1.44
11	73.75	1.05	0.25	0.14	1.44
12	63.35	1.16	0.19	0.15	1.50
13	67.85	1.00	0.22	0.16	1.38
14	63.45	1.03	0.21	0.11	1.35
15	66.75	1.05	0.2	0.14	1.39
16	65.65	1.00	0.21	0.15	1.36
17	79.75	1.01	0.17	0.17	1.35
18	56.35	1.00	0.27	0.17	1.44
Overall mean $\pm$ SE	64.16 $\pm$ 1.35	1.04 $\pm$ 0.01	0.20 $\pm$ 0.01	0.15 $\pm$ 0.01	1.38 $\pm$ 0.01

**Table 06: Average daily nutrient availability to Murrah buffalo calves**

Animal no.	Body weight (kg)	DM intake (kg)	CP (g)	TDN (g)	Ca (g)	P (g)	Fe (mg)	Cu (mg)	Zn (mg)	Mn (mg)	Co (mg)
1	59.95	1.33	56.62	572.76	6.12	2.56	492.1	9.97	10.86	62.82	0.11
2	62.53	1.33	57.34	572.68	6.08	2.56	498.06	9.39	10.56	62.23	0.15
3	61.67	1.30	58.43	570.64	6.63	2.56	460.00	9.87	10.95	62.17	0.16
4	62.75	1.34	58.88	570.64	6.75	2.98	468.06	9.67	10.78	62.17	0.12
5	58.87	1.47	58.49	570.94	6.85	2.89	486.03	9.98	10.81	62.67	0.12
6	61.75	1.39	57.64	570.74	5.98	2.67	492.07	8.85	10.89	62.06	0.08
7	62.90	1.34	58.32	571.94	6.63	2.45	484.01	8.86	10.96	62.56	0.04
8	65.75	1.35	58.86	570.54	5.95	2.89	485.56	9.67	10.87	62.13	0.36
9	55.87	1.39	57.88	572.83	6.34	2.83	499.12	9.89	10.63	62.72	0.13
10	65.85	1.44	58.67	570.84	6.45	2.65	472.06	10.79	10.89	62.87	0.12
11	73.75	1.44	59.89	570.83	6.64	2.75	591.03	9.98	10.71	62.87	0.06
12	63.35	1.50	58.45	571.77	6.46	2.83	496.01	10.62	10.92	62.11	0.03
13	67.85	1.38	58.96	572.82	6.22	2.45	596.11	9.45	10.84	62.47	0.07
14	63.45	1.35	56.98	572.57	6.81	2.34	489.08	9.54	10.72	62.32	0.09
15	66.75	1.39	58.73	570.74	6.84	2.98	589.04	10.89	10.77	62.86	0.14
16	65.65	1.36	58.83	571.94	6.75	2.56	587.11	10.95	10.74	62.11	0.12
17	79.75	1.35	59.67	572.94	6.76	2.22	594.23	9.67	10.82	62.84	0.12
18	56.35	1.44	56.34	569.74	5.87	2.56	492.05	9.97	10.86	62.31	0.11
Overall mean $\pm$ SE	64.16 $\pm$ 1.35	1.38 $\pm$ 0.01	58.26 $\pm$ 0.23	571.55 $\pm$ 0.24	6.45 $\pm$ .08	2.65 $\pm$ 0.05	515.10 $\pm$ 11.74	9.89 $\pm$ 0.14	10.81 $\pm$ 0.03	62.46 $\pm$ 0.07	0.12 $\pm$ 0.02

**Table 7: Average daily requirements and availability of nutrients in Murrah buffalo calves**

	Body weight (kg)	DM (kg)	CP (g)	TDN (g)	Ca (g)	P (g)	Fe (mg)	Cu (mg)	Zn (mg)	Mn (mg)	Co (mg)
<b>Requirements (ICAR,2013)</b>	70	1.8	287	1140	5.1	4.5	500	9.00	24.00	42.00	0.60
<b>Availability (Per animal per day)</b>	64.16	1.3	58.26	571.55	6.45	2.65	515.10	9.89	10.81	62.46	0.12
<b>Excess /Deficient</b>			(-)228.74	(-)568.45	(+)1.35	(-)1.85	(+)15.10	(+)0.89	(-)13.20	(+)20.46	(-)0.48
<b>Excess/Deficient%</b>			(-)79.70	(-)49.86	(+)26.47	(-)41.11	(+)3.02	(+)9.83	(-)54.96	(+)0.48	(-)79.5

Excess (+), Deficient (-)



### Nutrient availability

The average daily nutrient intake and body weight of Murrah buffalo calves are summarized in Table 06. The average body weight of the calves was  $64.16 \pm 1.35$  kg, with an average dry matter (DM) intake of  $1.38 \pm 0.01$  kg/day. The calcium and phosphorus intake averaged  $6.45 \pm 0.08$  g and  $2.65 \pm 0.05$  g, respectively. Among the trace minerals, iron intake was the highest at  $515.10 \pm 11.74$  mg, followed by manganese at  $62.46 \pm 0.07$  mg, zinc at  $10.81 \pm 0.03$  mg, copper at  $9.89 \pm 0.14$  mg and cobalt at  $0.12 \pm 0.02$  mg. These values indicated the overall nutrient availability from the feed ration provided to the calves.

### Nutritional deficiency/excess

The nutritional deficiency/excess data for minerals in a 64.16 kg (as analysed and presented in Table 09) body weight is presented in Table 08. The calves exhibited a deficiency of CP (228.74g, 79.70%), TDN,

(568.45kg, 49.86%), P(1.85g, 41.11%), Co (0.48mg, 79.5%) and Zn (13.20mg, 54.96%). Conversely, there was an excess of Ca (1.35 g, 26.47%), Fe(15.20mg, 3.20%), Cu (0.89 mg, 9.83%), and Mn(20.46 mg, 0.48%).

### Effect of strategic nutrient supplementation on dry matter intake (kg/d) of Murrah buffalo calves

The fortnightly feed intake (kg) of animals across the experimental groups is summarized in Table 08. At the beginning (0 fortnight), feed intake(kg/d) in the control group ( $T_0$ ) was  $2.46 \pm 0.03$ , while treatment groups  $T_1$  and  $T_2$  showed significantly higher values of  $2.52 \pm 0.05$  and  $2.65 \pm 0.06$ , respectively ( $p \leq 0.05$ ). The average dry matter intake (kg/d) in control ( $T_0$ ) and treatment ( $T_1$  and  $T_2$ ) groups during the 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup> fortnights were  $2.70 \pm 0.03$ ,  $2.75 \pm 0.06$  and  $2.81 \pm 0.06$ ;  $3.08 \pm 0.06$ ,  $2.92 \pm 0.05$  and  $3.02 \pm 0.04$ ;  $3.32 \pm$



0.05,  $3.21 \pm 0.07$  and  $3.41 \pm 0.06$ ;  $3.43 \pm 0.06$ ,  $3.48 \pm 0.08$  and  $3.64 \pm 0.08$ . In 5<sup>th</sup> fortnight, the average feed intake (kg/d) recorded was  $3.27 \pm 0.06$  in  $T_0$ ,  $3.76 \pm 0.10$  in  $T_1$  and  $3.83 \pm 0.09$  kg in  $T_2$ , with  $p \leq 0.01$ ) indicating a higher significant difference. In the 6<sup>th</sup> fortnight,  $T_0$  had an average feed intake(kg/d) of  $3.42 \pm 0.05$ , while  $T_1$  and  $T_2$  showed higher feed intake of  $3.95 \pm 0.10$  and  $4.04 \pm 0.10$ , respectively, with ( $p \leq 0.01$ ).

The overall mean feed intake (kg/d) was  $3.10 \pm 0.03$  for  $T_0$ , compared to significantly higher averages of  $3.23 \pm 0.07$  ( $T_1$ ) and  $3.34 \pm 0.06$  ( $T_2$ ). In the present study, significantly increased DMI was observed in the treatment groups in comparison to the control group.

The current study's findings are consistent with those of Zhou *et al.* (2015), who reported higher intakes of DM in cows fed on a high-energy diet than on low low-energy feed diet. Similarly, Hailu *et al.* (2011) and Tufarelli *et al.* (2011) also

observed higher DM intake in sheep fed higher levels of concentrate. Wang *et al.* (2019) reported that lambs' diet supplemented with 0.5 kg extra concentrate per day had significantly higher DM intake than those fed on the control diet. Furthermore, several other reports suggest that DM intake was significantly increased in buffaloes and sheep fed concentrate with an elevated crude protein (CP) content. Guire *et al.* (2013), Sweeny *et al.* (2014), Kang *et al.* (2015).

The probable explanation for the higher DM intake owing to the fact that, the customized supplement has sufficient fermentable energy and trace minerals to support and create a conducive environment for the growth of rumen microbes, leading to optimized rumen fermentation. Microbial growth and dry matter intake (DMI) were found to be strongly positively correlated (Seo *et al.*, 2013 and Uddin *et al.*, 2015). Similar results were also



reported by Chandra *et al.* (2015), Ojha *et al.* (2015) and Mahfuz *et al.* (2018).

In contrast to current findings, Kalita *et al.* (2010) discovered that there was no significant variation in dry matter intake between the two groups in nondescript male calves. Group I and Group II were fed a restricted amount of concentrate ration with and without mineral mixture at a 2% level, respectively. According to Sharma *et al.* (2011), there was no discernible variation in the overall dry matter consumption of the various animal groups. Likewise, Nagabhushana *et al.* (2008) reported that dry matter intake in crossbred calves showed no significant variation among the treatment groups. Similarly, Mohapatra *et al.* (2012) found that supplementing cows with an area-specific mineral mixture had no significant effect on their dry matter consumption. Similarly, Mishra *et al.* (2016) found no significant difference ( $p > 0.05$ ) in dry matter

(DM) intake between the dietary regimens. According to Hassan *et al.* (2016), there was no discernible variation in the total amount of feed consumed by the several groups of growing buffalo calves.

Contrary to our findings, no significant difference in dry matter intake between the treatment and control groups was documented by various authors Kumar *et al.* (2015), Sahoo *et al.* (2017) and Dixit *et al.* (2021).

#### **Growth performance of Murrah buffalo calves and effect of strategic nutrient supplementation on the body weight (kg) of Murrah buffalo calves**

The fortnightly changes in body weight of calves across the experimental groups are illustrated in Table 09. At the beginning of the study (0 fortnight), the body weight (kg) in the control group ( $T_0$ ) was  $62.79 \pm 6.07$ , while the treatment groups  $T_1$  and  $T_2$  had initial weights of  $62.23 \pm 4.77$  and  $62.66 \pm 2.45$ , respectively, the statistical analysis



specifies no significant differences ( $p > 0.05$ ) among the groups.

As the experiment progressed, body weight consistently increased across all groups. By the 1<sup>st</sup> fortnight, body weight (kg) reached  $66.11 \pm 6.16$  in  $T_0$ ,  $66.59 \pm 4.99$  in  $T_1$ , and  $67.41 \pm 2.45$  in  $T_2$ . The body weight (kg) changes in control ( $T_0$ ) and treatment ( $T_1$  and  $T_2$ ) groups during the 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> fortnights were  $69.85 \pm 6.26$ ,  $71.73 \pm 5.12$ , and  $73.07 \pm 2.50$ ;  $73.82 \pm 6.35$ ,  $78.12 \pm 5.06$ , and  $80.05 \pm 2.57$ ;  $77.45 \pm 6.41$ ,  $84.47 \pm 4.79$  and  $87.81 \pm 2.51$ ;  $81.1 \pm 6.34$ ,  $91.86 \pm 4.8$  and  $96.1 \pm 2.52$ , respectively. At the final measurement in the 6<sup>th</sup> fortnight,  $T_0$  calves weighed(kg) less  $85.03 \pm 6.31$ , as compared to  $99.55 \pm 4.79$  in  $T_1$  and  $104.54 \pm 2.60$  in  $T_2$ , with treatment groups showing significantly higher values than the control ( $p \leq 0.05$ ).

Overall, the mean body weight (kg) throughout the study was  $73.74 \pm 6.27$  for the control group ( $T_0$ ),  $79.22 \pm 4.9$  for  $T_1$ , and  $81.67 \pm 2.5$  for

$T_2$ . While  $T_1$  and  $T_2$  showed higher average body weights than  $T_0$ , these differences were not statistically significant ( $p > 0.05$ ) across the experimental period. Though the difference in average body weight was not significant, groups  $T_1$  and  $T_2$  exhibited 7.43% and 10.75% higher growth, respectively.

Similar to our findings, Nagabhushana *et al.* (2008) reported no discernible variation in body weight between treatment groups. The control group received a basal diet of wheat straw *ad libitum* with a concentrate mixture, without cobalt or mineral supplementation. The other two groups received the same basal diet and concentrate but were supplemented with cobaltous chloride at 1 ppm cobalt and 6 ppm cobalt, respectively. Likewise, Khan *et al.* (2015), reported that differences in body weight at all stages were no significant. Singh *et al.* (2015), dietary protein supplementation was found to have



no significant impact on body weight in buffalo heifers. In crossbreed calves, similarly, Nagalakshmi *et al.* (2018), supplementation of (80 ppm Zn) in the control group and (60 ppm Zn) in the experimental group had comparable between the dietary groups.

Contrary to our finding, Tewari *et al.* (2014) concluded that calves' body weight changes were considerably ( $p < 0.01$ ) substantial in the treatment group than in the control group. Similarly, Hassan *et al.* (2016) found that calves fed a supplemented zinc sulphate or zinc methionine feed had significantly higher body weights ( $p < 0.05$ ) than those fed a control ration Gouda *et al.* (2017) found that supplemented groups had significantly ( $p \leq 0.05$ ) higher body weights than the control group when the later were fed ASMM @ 50 g per day while the former were kept in accordance with the farmer's traditional practices (straw-based diet with

locally available concentrate) without any nutritional supplementation.

### **Effect of strategic nutrient supplementation on average daily gain (g/d) of Murrah buffalo calves**

The fortnightly average daily gain (ADG) in  $T_0$ ,  $T_1$ , and  $T_2$  groups is presented in Table 11. In the first fortnight, the average daily gain (g) for the control group ( $T_0$ ) was  $221 \pm 10.31$ , while in the treatment groups  $T_1$  and  $T_2$ , the ADG values were significantly higher at  $290.78 \pm 15.99$  and  $316.78 \pm 15.83$ , respectively ( $p \leq 0.01$ ). This trend of higher ADG (g) in the treatment groups continued in the 2<sup>nd</sup> fortnight, where  $T_0$  recorded an ADG of  $250.34 \pm 11.33$ , whereas  $T_1$  and  $T_2$  showed significantly higher gains of  $343.12 \pm 15.6$  and  $377.67 \pm 11.53$ , respectively ( $p < 0.01$ ). Weight gain (g) in  $T_0$ ,  $T_1$ , and  $T_2$  during the 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup>, and 6<sup>th</sup> fortnights were  $264.11 \pm 21.67$ ,  $425.78 \pm 36.48$  and  $464.92 \pm 14.30$ ;  $241.67 \pm 7.67$ ,  $443.78 \pm 28.27$  and





517.08  $\pm$  10.25; 243.45  $\pm$  17.25, 492.78  $\pm$  11.58 and 553.00  $\pm$  18.03; 261.89  $\pm$  9.35, 512.78  $\pm$  12.37 and 562.23  $\pm$  13.8. The overall mean average daily gain across the study period was 247.08  $\pm$  4.75 for T<sub>0</sub>, 418.17  $\pm$  9.4 for T<sub>1</sub>, and 465.28  $\pm$  6.81 for T<sub>2</sub>, indicating significant differences among the groups ( $p \leq 0.01$ ). ADG (g) was considerably ( $P < 0.05$ ) higher in treatment groups than in the control group, despite the fact that overall body weight changes did not differ significantly across groups because of considerable differences within the group.

Consistent with the present study, Meyer *et al.* (2010) reported a significant ( $P < 0.05$ ) improvement in average daily gain among supplemented groups compared to the control group. When 0.5 g of area-specific mineral mixture was supplemented. Similarly, when given a basal diet in the control group and supplemented with Fe, Mn, Zn, Cu, I, Co, Se, vitamin-A,

vitamin-D, and vitamin-E in the treatment group, Mishra *et al.* (2016) observed a substantial ( $p < 0.01$ ) increase in average daily gain in the T group compared to the control (C) group. According to Gouda *et al.* (2017), the average daily gain of the supplemented groups was significantly ( $p \leq 0.05$ ) higher than that of the control group when the latter were fed ASMM @ 50 g per day per animal while the former were kept in accordance with the farmer's traditional practices (straw-based diet with locally available concentrate) without any nutritional supplementation. Similarly, according to Sahoo *et al.* (2016), there was a significant difference ( $p < 0.05$ ) was observed in the average daily gain (g) between all treatment groups and the control group. In this study, the control group (C) was managed according to traditional farmer practices, while the treatment groups received additional



supplementation. The T1 group was supplemented with 50 g of mineral mixture per day per animal, and the T2 group was given 100 g of bypass fat per animal per day in combination with 50 g of mineral mixture. Similarly, Hassan *et al.* (2016) found that calves fed a supplemented zinc-methionine or zinc sulphate ration had significantly higher average daily gain ( $p \leq 0.05$ ) than those fed a control ration. This was because the supplemented zinc-methionine ration increased methionine absorption and stimulated the activities of enzymes involved in nutrient digestibility, feeding value, and feed efficiency. Similarly, Rojita Yengkhom *et al.* (2018) found that the commercial and booster-mineral supplement group's average daily increased during the 135-day period was considerably higher ( $p < 0.01$ ) than the control groups in kid. Chaudhary *et al.* (2024) reviewed that the average daily gain was markedly higher ( $p \leq 0.01$ ) in the novel feed

supplement group compared to the basal roughages and concentrate mixture 20% of DMI as per the farmers' practices (control group).

In contrast to our results, Mandal *et al.* (2008) found no discernible variation in the body weight growth of crossbred calves across groups. In crossbred calves of different groups, Sharma *et al.* (2010) showed no significant difference in average daily weight growth; for G 1, G 2, and G 3, the average daily weight gain was 509, 556, and 496 g, respectively. According to Meher *et al.* (2017), there was no discernible difference ( $P > 0.05$ ) in the crossbred cows' average daily weight gain (g) between the control and treatment groups. In this study, the control group (T0) followed the farmer's traditional practices without additional nutritional intervention. In the T1 group, animals were supplemented daily with 50 g of an ASMM per animal. In the T2 group, animals also received this mineral mixture, and, in addition, each



animal was administered an injection of fat-soluble vitamins (Vitamin A, D3, and E) on both day 0 and day 7 of the experiment. Similarly, Mudgal *et al.* (2008) revealed that the average daily gain was found to be similar ( $p > 0.05$ ) among the four groups. Contrary to our findings, a nonsignificant difference in average daily gain between the treatment and control groups was reported by various authors, Lammer and Heinrichs *et al.* (2000), Hadiya *et al.* (2009) and Das *et al.* (2014).

#### **Effect of strategic nutrient supplementation on the feed conversion ratio of Murrah buffalo calves**

The feed conversion ratio (FCR) at fortnightly intervals is presented in Table 12. The FCR in the first fortnight for the control and treatment groups was  $12.33 \pm 0.57$ ,  $9.58 \pm 0.54$  and  $8.99 \pm 0.53$  in groups  $T_0$ ,  $T_1$ , and  $T_2$ , respectively, and it was statistically significant ( $P \leq 0.01$ ) between the groups. FCR in  $T_0$ ,  $T_1$

and  $T_2$  during the 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup>, and 6<sup>th</sup> fortnights were  $12.49 \pm 0.72$ ,  $8.6 \pm 0.44$  and  $8.06 \pm 0.28$ ;  $13.02 \pm 1.15$  for  $T_0$ ,  $7.81 \pm 0.70$  for  $T_1$ , and  $7.35 \pm 0.25$ ;  $14.26 \pm 0.49$ ,  $7.96 \pm 0.48$  and  $7.04 \pm 0.18$ ;  $13.67 \pm 0.78$ ,  $7.63 \pm 0.19$  and  $6.95 \pm 0.21$ ;  $13.14 \pm 0.53$ ,  $7.74 \pm 0.31$  and  $7.19 \pm 0.09$  for  $T_2$ , with statistically significant differences ( $p \leq 0.01$ ) between the control and treatment groups.

The overall average FCR across all fortnights was  $13.15 \pm 0.32$  for  $T_0$ ,  $8.22 \pm 0.2$  for  $T_1$ , and  $7.6 \pm 0.15$  for  $T_2$ , with the treatment groups ( $T_1$  and  $T_2$ ) demonstrating significantly improved feed conversion efficiency compared to the control group ( $p \leq 0.01$ ). These findings indicate that nutrient supplementation enhanced the feed conversion ratio in the treatment groups, resulting in a more efficient utilization of feed.

According to Sawant *et al.* (2013), the feed conversion efficiency was lowest ( $p < 0.05$ ) in the control group and highest ( $p \leq 0.05$ ) in the



heifers of group T3, followed by T2. The control group's heifers were given 1.0 kg of concentrate, 5 kg of green fodder, and Jowar kadbi at will. The same was given to the treatment group (T2), along with an extra 30 g of mineral mixture; the heifers in the treatment group (T3) also received 30 g of minerals mixed with vitamins. Similarly, Hassan *et al.* (2016) showed that supplementing growing buffalo calves with zinc methionine or zinc sulphate increased feed conversion efficiency significantly ( $p < 0.05$ ) when compared to a control diet devoid of supplements.

In contrast, earlier research by Das *et al.* (2014), concluded that the feed conversion ratio was found to be similar in all three groups. Similarly, Mishra *et al.* (2016) studied that feed conversion

efficiency did not differ significantly between groups when fed a basal diet in the control group and supplemented with Fe, Mn, Zn, Cu, I, Co, Se, vitamin-A, vitamin-D, vitamin-E in the treatment group.

Contrary results were also summarized by various authors, including Basra *et al.* (2003), Chaturvedi *et al.* (2009), and Singh *et al.* (2015), who reported a non-significant difference in feed conversion ratio between the treatment and control groups.

### Conclusions

Dietary supplementation with strategic nutrients significantly improved body weight, dry matter intake, average daily gain, and feed conversion ratio in the treatment groups.



**Table 08: Effect of strategic nutrient supplementation on dry matter intake (kg/d) of Murrah buffalo calves**

Fortnights	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	p value
0	2.46 <sup>a</sup> ± 0.03	2.52 <sup>ab</sup> ± 0.05	2.65 <sup>b</sup> ± 0.06	0.03
1	2.70 ± 0.03	2.75 ± 0.06	2.81 ± 0.06	0.29
2	3.08 ± 0.06	2.92 ± 0.05	3.02 ± 0.04	0.11
3	3.32 ± 0.05	3.21 ± 0.07	3.41 ± 0.06	0.07
4	3.43 ± 0.06	3.48 ± 0.08	3.64 ± 0.08	0.11
5	3.27 <sup>a</sup> ± 0.06	3.76 <sup>b</sup> ± 0.1	3.83 <sup>b</sup> ± 0.09	0.01
6	3.42 <sup>a</sup> ± 0.05	3.95 <sup>b</sup> ± 0.1	4.04 <sup>b</sup> ± 0.10	0.01
Overall mean ± SE	3.10 <sup>a</sup> ± 0.03	3.23 <sup>ab</sup> ± 0.07	3.34 <sup>b</sup> ± 0.06	0.01

Means bearing different superscripts in a row differ significantly ( $p \leq 0.05$ )

Means bearing different superscripts in a row differ significantly ( $p \leq 0.01$ )

**Table 09: Effect of strategic nutrient supplementation on body weight (kg) of Murrah buffalo calves**

Fortnights	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	P value
0	62.79 ± 6.07	62.23 ± 4.77	62.66 ± 2.45	1.00
1	66.11 ± 6.16	66.59 ± 4.99	67.41 ± 2.45	0.98
2	69.85 ± 6.26	71.73 ± 5.12	73.07 ± 2.50	0.90
3	73.82 ± 6.35	78.12 ± 5.06	80.05 ± 2.57	0.66
4	77.45 ± 6.41	84.47 ± 4.79	87.81 ± 2.51	0.33
5	81.10 ± 6.34	91.86 ± 4.80	96.10 ± 2.52	0.11
6	85.03 <sup>a</sup> ± 6.31	99.55 <sup>b</sup> ± 4.79	104.54 <sup>b</sup> ± 2.60	0.03
Overall mean ± SE	73.74 ± 6.27	79.22 ± 4.90	81.67 ± 2.50	0.51
Percent increase (%)		7.43	10.75	

Mean bearing different superscripts in a row differ significantly ( $p \leq 0.05$ )



**Table 11: Effect of strategic nutrient supplementation on average daily gain (g/d) of Murrah buffalo calves**

Fortnights	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	p value
1	221.00 <sup>a</sup> ± 10.31	290.78 <sup>b</sup> ± 15.99	316.78 <sup>b</sup> ± 15.83	0.01
2	250.34 <sup>a</sup> ± 11.33	343.12 <sup>b</sup> ± 15.60	377.67 <sup>b</sup> ± 11.53	0.01
3	264.11 <sup>a</sup> ± 21.68	425.78 <sup>b</sup> ± 36.48	464.92 <sup>b</sup> ± 14.30	0.01
4	241.67 <sup>a</sup> ± 07.67	443.78 <sup>b</sup> ± 28.27	517.08 <sup>c</sup> ± 10.25	0.01
5	243.45 <sup>a</sup> ± 17.25	492.78 <sup>b</sup> ± 11.58	553.00 <sup>c</sup> ± 18.03	0.01
6	261.89 <sup>a</sup> ± 09.35	512.78 <sup>b</sup> ± 12.37	562.20 <sup>c</sup> ± 13.80	0.01
Overall mean ± SE	247.08 <sup>a</sup> ± 04.75	418.17 <sup>b</sup> ± 09.40	465.28 <sup>c</sup> ± 06.81	0.01

Mean bearing different superscripts in a row differ significantly ( $p \leq 0.01$ )

**Table 12: Effect of strategic nutrient supplementation on the feed conversion ratio of Murrah buffalo calves**

Fortnights	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	p value
1	12.33 <sup>a</sup> ± 0.57	9.58 <sup>a</sup> ± 0.54	8.99 <sup>b</sup> ± 0.53	0.01
2	12.49 <sup>a</sup> ± 0.72	8.60 <sup>a</sup> ± 0.44	8.06 <sup>b</sup> ± 0.28	0.01
3	13.02 <sup>a</sup> ± 1.15	7.81 <sup>a</sup> ± 0.70	7.35 <sup>b</sup> ± 0.25	0.01
4	14.26 <sup>a</sup> ± 0.49	7.96 <sup>a</sup> ± 0.48	7.04 <sup>b</sup> ± 0.18	0.01
5	13.67 <sup>a</sup> ± 0.78	7.63 <sup>a</sup> ± 0.19	6.95 <sup>b</sup> ± 0.21	0.01
6	13.14 <sup>a</sup> ± 0.53	7.74 <sup>a</sup> ± 0.31	7.19 <sup>b</sup> ± 0.09	0.01
Overall mean ± SE	13.15 <sup>a</sup> ± 0.32	8.22 <sup>a</sup> ± 0.20	7.60 <sup>b</sup> ± 0.15	0.01

Mean bearing different superscripts in a row differs significantly ( $p \leq 0.01$ )

## References

Banerjee, G. C. (1998). A textbook of animal husbandry 8th Edn., Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi, pp 778.

Basra, M. J., Khan, M. A., Nisa, M., Riaz, M., Tuqeer, N. A. and Saeed, M. N. (2003). Nili-Ravi buffalo I. Energy and protein requirements of 6-9 months old calves. *International Journal of Agriculture and Biology*, 5(3): 377-379.



- Chandra, G., Aggarwal, A., Singh, A. K. and Kumar, M. (2015). Effect of vitamin E and zinc supplementation on milk yield, milk composition, and udder health in Sahiwal cows. *Animal Nutrition and Feed Technology*, **15**(1): 67-78.
- Chaturvedi, O. H., Kumar, S., Mishra, A. K., Arora, A. L. and Karim, S. A. (2009). Effect of complete feed or grazing and supplementation of lambs on performance, nutrient utilization and feed cost of production. *The Indian Journal of Animal Sciences*, **79**(9): 917-920.
- Chaudhary, S. K., Dutta, N., Jadhav, S. E., Singh, G., Singh, S. K. and Tewari, D. (2024). Effect of strategic supplementation of a novel feed supplement on zoo-technical attributes, metabolic profile, production and reproduction performance of buffaloes. *Buffalo Bulletin*, **43**(2): 209-225.
- Das, L. K., Kundu, S. S., Datt, C. and Kumar, D. (2014). Metabolizable protein requirements of Sahiwal calves fed on berseem (*Trifolium alexandrinum*) fodder based rations. *Indian Journal of Animal Nutrition*, **31**(1): 8-13.
- Dixit, S., Keshri, A., Vinay, V. V. and Kundu, S. S. (2021). Effect of graded levels of dietary crude protein on the enteric methane emissions in growing Murrah buffalo calves. *Indian Journal of Dairy Science*, **74** (6): 550-553.
- Gouda, G., Sethy, K., Swain, R. K., Mishra, S. K., Behera, K., Debata, N. R. and Mishra, B. (2017). Effect of area specific mineral mixture and hormonal interventions on growth, blood chemistry and mineral status of the reproductive disorders cross bred cattle. *Journal of Pharmaceutical Innovation*, **6**: 284-286.
- Hadiya, K. Maini, S., Rekhe, D.S. and Ravikanth, K. (2009). Accelerated growth programme with polyherbal formulations for dairy calves. *Veterinary World*, **2**(2): 62-64.
- Hailu, A., Melaku, S., Tamir, B. and Tassew, A. (2011). Body weight and carcass characteristics of Washera sheep fed urea treated rice straw supplemented with graded levels of concentrate mix. *Livestock Research for Rural Development*, **23**(8): 164-185.
- Hassan, E. H., Farghaly, M. M. and Solouma, G. M. (2016). Effect of zinc supplementation from inorganic and organic sources on nutrient digestibility, some blood metabolites and growth performance of growing buffalo calves. *Egyptian Journal of Nutrition and Feeds*, **19**(1): 37-46.
- ICAR (2013). Nutrient Requirements of Animals – Cattle and Buffalo. Indian Council of Agricultural Research, New Delhi.
- Kalita, D. J., Sarma, B. C. and Sarmah, B. K. (2010). Effect of mineral supplementation on growth, retention percentage and serum mineral profile of calves. *Indian Journal of Animal Research*, **44**(3): 208-210.
- Kang, S., Wanapat, M., Phesatcha, K. and Norrapoke, T. (2015). Effect of protein level and urea in concentrate mixture on feed intake and rumen fermentation in swamp buffaloes fed rice straw-based diet. *Tropical Animal Health and Production*, **47**(4): 671-679.



- Khan, H. M., Mohanty, T. K., Bhakat, M., Gupta, A. K. and Mondal, G. (2015). Effect of vitamin E and mineral supplementation during peri-partum period on BCS, body weight and calf performance in Murrah buffaloes. *Buffalo Bulletin*, **34**(1): 79-85.
- Kumar, S., Sehgal, J. P., Puniya, A. K. and Kumari, R. (2015). Growth performance and fibre utilization of Murrah male buffalo calves fed wheat straw based complete feed blocks incorporated with superior anaerobic fungal zoospores (*Neocallimastix* sp.GR-1). *The Indian Journal of Animal Sciences*, **85**(3): 275-81.
- Lammers, B. P. and Heinrichs, A. J. (2000). The response of altering the ratio of dietary protein to energy on growth, feed efficiency, and mammary development in rapidly growing prepubertal heifers. *Journal of Dairy Science*, **83**(5): 977-983.
- Mahfuz, S. U., Islam, M., Chowdhury, M. R., Islam, S., Hasan, M. K. and Uddin, M. N. (2018). Influence of concentrate supplementation on production and reproduction performance of female Black Bengal goat. *Indian Journal of Animal Research*, **52**(5): 735-739.
- Mandal, G. P., Dass, R. S., Garg, A. K., Varshney, V. P. and Mondal, A. B. (2008). Effect of zinc supplementation from inorganic and organic sources on growth and blood biochemical profile in crossbred calves. *Journal of Animal and Feed Sciences*, **17**(2): 147-156.
- McGuire, D.L., Bohnert, D.W., Schauer, C.S., Falck, S.J. and Cooke, R.F. (2013). Daily and alternate day supplementation of urea or soybean meal to ruminants consuming low-quality cool-season forage: I-Effects on efficiency of nitrogen use and nutrient digestion. *Livestock Science*, **155**(2-3): 205-213.
- Mishra, A., Singh, P., Verma, A. K. and Ojha, B. K. (2016). Effect of micronutrients supplement on nutrient utilization and growth performance in pre-ruminant calves. *Journal of Animal Research*, **6**(2): 251-255.
- Mohapatra, P., Swain, R. K., Mishra, S. K., Sahoo, G. and Rout, K. K. (2012). Effect of supplementation of area specific mineral mixture on reproductive performance of the cows. *The Indian Journal of Animal Sciences*, **82**(12): 1558-1563.
- Mudgal, V., Garg, A. K. and Dass, R. S. (2008). Effect of zinc, copper and selenium supplementation on growth rate and nutrient utilization in Buffalo (*Bubalus bubalis*) calves. *Indian Journal of Animal Nutrition*, **25**(3): 272-277.
- Nagabhushana, V., Sharma, K., Pattanaik, A. K. and Dutta, N. (2008). Effect of cobalt supplementation on performance of growing calves. *Veterinary World*, **1**(10): 299-302.
- Ojha, B. K., Dutta, N., Pattanaik, A. K., Singh, S. K. and Narang, A. (2015). Effect of pre-partum strategic supplementation of concentrates on colostrum quality and performance of buffalo calves. *Animal Nutrition and Feed Technology*, **15**(1): 41-49.
- Rojita Yengkhom, R. Y., Verma, A. K., Narayan Dutta, N. D., Jadhav, S. E. and Pattanaik, A. K. (2018). Effect of a customized mineral supplement





- on nutrient metabolism, serum mineral profile and growth performance of kids. *Animal Nutrition and Feed Technology*, 18(2): 177-187.
- Sahoo, B., Kumar, R., Garg, A. K., Mohanta, R. K., Agarwal, A. and Sharma, A. K. (2017). Effect of supplementing area specific mineral mixture on productive performance of crossbred cows. *Indian Journal of Animal Nutrition*, 34(4): 414-419.
- Sawant, D. N., Todkar, S. R. and Sawant, P. J. (2013). Effect of supplementation of minerals and vitamins on growth performance of indigenous heifers. *Indian Journal of Animal Nutrition*, 30(4): 387-391.
- Seo, J.K., Kim, M.H., Yang, J.Y., Kim, H.J., Lee, C.H., Kim, K.H. and Ha, J.K. (2013). Effects of synchronicity of carbohydrate and protein degradation on rumen fermentation characteristics and microbial protein synthesis. *Asian-Australasian Journal of Animal Science*, 26: 358-365.
- Sharma, D., Tiwari, D. P. and Mondal, B. C. (2010). Performance of the crossbred female calves fed complete ration as mash or block vis-a-vis conventional ration. *The Indian Journal of Animal Sciences*, 80(6): 556-560.
- Sharma, J., Kumar, A., Tiwari, D. P. and Mondal, B. C. (2011). Effect of dietary supplementation of calcium, copper and manganese on nutrient utilization, growth, blood biochemical and mineral profile in crossbred heifers. *The Indian Journal of Animal Sciences*, 81(5): 493-497.
- Singh, S., Kushwaha, B. P., Maity, S. B., Singh, K. K. and Das, N. (2015). Effect of dietary protein on intake, nutrients utilization, nitrogen balance, blood metabolites, growth and puberty in growing Bhadawari buffalo (*Bubalus bubalis*) heifers. *Tropical Animal Health and Production*, 47: 213-220.
- Snedecor, G. W. and Cochran, W. G. (1994). *Statistical methods*, Iowa State University press, Ames. Iowa, USA.
- Sweeny, J.P., Surridge, V., Humphry, P.S., Pugh, H. and Mamo, K. (2014). Benefits of different urea supplementation methods on the production performances of Merino sheep. *Veterinary Journal*, 200(3): 398-403.
- Tewari, D., Jain, R. K. and Mudgal, V. (2014). Effect of strategic nutrient supplementation on the reproductive performance of anoestrus crossbred cattle in Malwa region of Madhya Pradesh. *Indian Journal of Animal Research*, 48(6): 580-584.
- Tufarelli, V., Lacalandra, G. M., Aiudi, G., Binetti, F. and Laudadio, V. (2011). Influence of feeding level on live body weight and semen characteristics of Sardinian rams reared under intensive conditions. *Tropical Animal Health and Production*, 43(2): 339-345.
- Uddin, M.J., Khandaker, Z.H., Khan, M. and Khan, M.M.H. (2015). Dynamics of microbial protein synthesis in the rumen-a review. *Annals of Veterinary and Animal Science*, 2(5): 116-131.
- Wang, C., Zhao, Y., Aubry, A., Arnott, G., Hou, F. and Yan, T. (2019). Effects of concentrate input on nutrient



---

utilization and methane  
emissions of two breeds of ewe  
lambs fed fresh ryegrass.

Translational Animal Science, 3:  
485-492.